

S/051/61/011/006/004/012
E032/E114

AUTHORS: Tatevskiy, M., Koptev, G.S., and Mal'tsev, A.A.

TITLE: Vibrational spectra of the B_2O_3 and B_2S_3 molecules

PERIODICAL: Optika i spektroskopiya, v.11, no.6, 1961, 724-729

TEXT: In a previous paper (A.A. Mal'tsev, V.N. Matveyev, V.M. Tatevskiy, DAN SSSR, v.137, 123, 1961) it was shown that the frequency of the antisymmetric BO vibration is about 900 cm^{-1} , while D. White, D.E. Mann, P.N. Walsh and A. Sommer (Ref.3, J.Chem Phys., v.32, 481, 1960) reported the figure of 1302 cm^{-1} . The aim of the present work was to explore the possible reasons for this discrepancy. The authors describe a calculation of the vibrational spectra of the $B_2^{10}O_3$ and $B_2^{11}O_3$ molecules, and have

recalculated the spectra of $B^{10}B^{11}O_3$, $B_2^{10}S_3$ and $B_2^{11}S_3$. The calculated frequencies of bands in the vibrational spectra of $B_2^{10}O_3$, $B^{10}B^{11}O_3$, $B_2^{11}O_3$, $B_2^{10}S_3$ and $B_2^{11}S_3$ are shown in the following table:

Card 1/ 3

Vibrational spectra of the B_2O_3 and

S/051/61/011/006/004/012
E032/E114

| Vibration | Symmetry | $B^{10}O_3$ | $B^{10}B^{11}O_3$ | $B^{11}O_3$ | $B^{10}S_3$ | $B^{11}S_3$ |
|--|----------------|-------------|-------------------|-------------|-------------|-------------|
| ν_1 ν_2 ν_3 ν_4 | A ₂ | 2084 | 2022 | 2015 | 1359 | 1306 |
| | | 844 | 829 | 815 | 425 | 409 |
| | | 755 | 754 | 749 | 399 | 396 |
| | | 161 | 161 | 160 | 66 | 64 |
| ν_5 ν_6 ν_7 | B ₁ | 2097 | 2091 | 2029 | 1364 | 1317 |
| | | 938 | 936 | 933 | 463 | 461 |
| | | 682 | 668 | 658 | 347 | 333 |
| ν_8 | A ₂ | 778 | 768 | 753 | 374 | 360 |
| ν_9 | B ₂ | 700 | 685 | 673 | 351 | 336 |



The calculations show that the bands 1302 and 742 cm^{-1} which are observed in the infrared emission spectrum of vapours over boric anhydride are due to the liquid phase and not the B_2O_3

Card 2/ 3

Vibrational spectra of the B_2O_3 and .. S/051/61/011/006/004/012
E032/E114

molecule. The 993 cm^{-1} band in the spectrum of boron sulphide is due to the spectrum of liquid B_2S_3 . It is suggested that in order to obtain more accurate values for the force constant and in order to increase the accuracy of the vibrational frequencies, more accurate spectroscopic studies are necessary in the region of 900 cm^{-1} where it is expected that there are bands due to antisymmetric BO vibrations.

There are 1 table and 17 references: 5 Soviet-bloc and 12 non-Soviet-bloc. The four most recent English language references read as follows:

Ref. 3: D. White, D.E. Mann, P.N. Walsh, A. Sommer, J.Chem.Phys., v.32, 481, 1960. ✓

Ref.11: A. Sommer, P.N. Walsh, D. White, J.Chem.Phys., v.33, 296, 1960.

Ref.12: C.W.F.T. Pistorius, J.Chem.Phys., v.31, 1454, 1959.

Ref.16: J.L. Parsons, M.E. Milberg. J.Amer.Ceram.Soc., v.43, 326, 1960.

SUBMITTED: January 2, 1961

Card 3/3

VYGODSKAYA, Ye.M.; LATEVSKIY, V.M.

Method of estimating the physicochemical properties of boron
alkyls. Vest. Mosk. un. Ser. 2: Khim. 19 no.6:8-10 N-D '64.
(MIRA 18:3)

1. Kafedra fizicheskoy khimii Moskovskogo universiteta.

KOPEV, G.S.; PANCHENKO, Yu.N.; TYULIN, V.I.; TATEVSKIY, V.M.

Calculating the frequency and vibration pattern of two isotopic forms of the molecules of 1,3-butadiene C_4H_6 and C_4D_6 . Opt. i spektr. 19 no.2:194-197 Ag '65. (MIRA 18:8)

GEY, G.; YAKOVLEV, S.S.; ZAKHAROV, V.S.

Dipole moment of components of the general formula in part 2.
Vest.Mosk.univ.Khim. 20 no.3:13-19. 1965.

(MIRA 18:2)

L. Moskovskiy universitet, kafedra fizicheskoy khimii.

TATEVSKIY, V.M.; SPIRIDONOV, V.P.

Remarks on G.V. Bykov's article "The terms electronegativity and electrophilic characteristic of atoms in molecules" and on "Electronegativity of atoms (atomic cores) in molecules." Part 1. Zhur. fiz. khim. 39 no.5:1284-1289 My '65.

Remarks on G.V. Bykov's articles "Term of electronegativity and electrophilic characteristic of atoms in molecules" and "Electronegativity of atoms (atomic cores) in molecules. Part 2. Ibid.:1291-1295 (MIRA 18:8)

1. Moskovskiy gosudarstvennyy universitet imeni M.V. Lomonosova.

GEY, E.; YAROVY, S.S.; TATEVSKIY, W.M.

Dipole moments of compounds of the general formula $A_n B_{2n+2}$.
West. Mosk. un. Ser. 2:Khim. 20 no.4:3-6 J1-Ag '65.

(MIRA 18:10)

1. Kafedra fizicheskoy khimii Moskovskogo gosudarstvennogo universiteta.

Tate ~~et al.~~ V. Vibrations of polyatomic molecules. Akad.

CA 2

The catalytic action of aluminosilicates. IV. Transformation of cyclohexene over activated clay. A. P. Nikolaev, V. M. Tatevskii, and A. V. Prosk. *J. Gen. Chem. (U.S.S.R.)* 13, 706-8(1940); cf. *C.A.* 39, 062'. - Passage of cyclohexene over activated clay at 260° results in formation of methylcyclopentane and cyclohexane in the ratio 10:1. Small amts. of unsatd. and aromatic hydrocarbons are formed and there is a 14% loss as gas or tar. H. M. Leicester

ASSOCIATED METALLURGICAL LITERATURE CLASSIFICATION

| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 60 | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70 | 71 | 72 | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 80 | 81 | 82 | 83 | 84 | 85 | 86 | 87 | 88 | 89 | 90 | 91 | 92 | 93 | 94 | 95 | 96 | 97 | 98 | 99 | 100 |
|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----|

PROCESSED AND PROPERTIES INDEX

3

Intensity of the combination scattering of halogen-substituted hydrocarbons. V. Tatarskii (State Univ., Moscow). J. Phys. Chem. (U.S.S.R.) 20, 129-31 (1946).—Raman spectra of mixts. of CCl₄ with another compd. were obtained, and the intensity I of some lines of the other compd. was compared with that of the 450 cm.⁻¹ line of CCl₄ taken for 100. The I was 180 for the CH₃I line 522 cm.⁻¹; 80, 72, and 71 for the CHCl₃ lines 261, 366, and 667; 225, 184, 124, and 78 for the CHBr₃ lines 164, 222, 539, and 655; 37 for the BrCl line 653; 85 for the Br₂ line 537; 171 for the BrI line 107; and 24 and 37 for the 653 and 733 lines of 1,2-dichloroethane. The I of propyl (or isopropyl) chloride, bromide, and iodide was detd. in other relative units. The spectrum of CH₃I could not be obtained because of photochem. decompa. The error was ±2 to ±7%. The results will be discussed later.

J. J. Bikerman

A S B - S L A METALLURGICAL LITERATURE CLASSIFICATION

| | | | |
|----------|----------|----------|----------|
| 12000 72 | 12000 72 | 12000 72 | 12000 72 |
| 12000 72 | 12000 72 | 12000 72 | 12000 72 |

TATEVSKIY, V. M.

Mar 1947

USSR/Chemistry - Ethylene
Chemistry - Ethane

"Calculation of the Reaction Equilibrium of Chlorine Substitution in Ethylene and Ethane," V. M. Tatevskiy, Izvestiya, Prof. A. V. Frost, 19 pp

"Vestnik Moskovskogo Universiteta" No 3
87-15-83
By methods of statistical thermodynamics from thermochemical and spectroscopic data, calculations have been made of the equilibrium constant and molar fractions in equilibrium mixture for 13 reactions of chlorine substitution in ethylene and ethane in the temperature range 0 - 600° C. The barrier between trans- and laevo-forms of dichloroethane has been approximately calculated, the value $V_0 = 3,000 - 2473$

Mar 1947

USSR/Chemistry - Ethylene (Contd)

4,000 cal/mol having been obtained, which agrees satisfactorily, within the limits of precision of the computation, with the results obtained by other methods.

2473

TATTOO V M

...

TATEVSKIY, V. M.

42039: ARONOV, O. L.; TATEVSKIY, V. M.; FROST, A. V. - Issledovanie potatsionnoy izomerii
hekotorykh I - Digalogenogtanov metodom kombinatsionnogo rasseyaniya sveta.
(Kratkoe sodержanie doklada). Izvestiya akad. Nauk SSSR, Seriya Fiz., 1948,
No. 5, S. 666.

SO: Letopis' Zhurnal'nykh Statey, Vol. 47, 1948.

TATEVSKIY, V. M.

25401. YURBEV, Y. K., TATEVSKIY, V.M. i GRAGEROV, I.P.
Spektry kombinatsionnogo rasseyaniya tiofana i ego gomologov. Zhurnal fiz.
Khimii, 1948, Vyp. 7., 783-86. Bibliogr: 6 Nazv. Sm Takzhe No. 25427

SO: Letopis' Zhurnal Statey, No. 30, Moscow, 1948

TATEVSKIY, V. M.

42097. TATEVSKIY, V. M. O kataliticheskoy aktivnosti sistemy adsorbent/
adsorbiruyemogo veshchestva. Vestnik Mosk. un-ta, 1948, No 8,
s. 77-83.- Bibliogr: 5 nazv.

SO: Letopis' Zhurnal'nykh Statey, Vol. 36, 1948

TATEVSKIY, V. M.

Tatevskiy, V. M. and Finkel'shteyn, A. I. "An approximate method of calculating determinants of a special type", Vestnik Mosk. un-ta, 1948, No. 10, p. 63-74.

SO: U-3042, 11 March 53, (Ietopis 'nykh Statey, No. 10, 1949).

TATEVSKIY, V.M.
ARONOV, O.L., TATEVSKIY, V.M., AND FROST, A.V.

"Study of Rotational Isomerism by the Combination Light Dispersion Method. IV. Spectra of Combination Dispersion of 1, 1., 2, 2-tetrachloroethane at Various Temperatures."
Vestnik Moskovskogo Universiteta, no. 11, 1948

TATEVSKIY, V. M.

Tatevskiy, V. M. - "Combination scattering spectra of aromatic C₈ hydrocarbons,"
Vestnik Mosk. un-ta, 1948, No. 11, p. 149-53

So: U- 3566, 15 March 53, (Letopis 'Zhurnal 'nykh Statey, No. 13, 1949)

TATEVSKIY, V. M.

USSR/Chemistry - Isomerism
Chemistry - Spectra

Sep/Oct 48

"Studies of the Rotational Isomerism of Some 1,2-Dihalogen-ethanes by the Method of Combined Dispersion of Light," D. L. Aronov, V.M. Tatevskiy, A. V. Frost, $\frac{1}{2}$ p

"Iz Ak Nauk SSSR, Ser Fiz" Vol XII, No 5

Studies combination dispersion spectra of 1,2-dichloroethane, 1-chloro, 2-bromoethane, 1,2-dibromoethane and 1,1,2,2-tetrachloroethane in the liquid phase over a wide range of temperatures. (Synopsis. For detailed description, see "Vest Moskov U" 1, 125, 1943.)

PA 19/49T4

TATEVSKIY, V. M.

USSR/Chemistry - Organic Compounds Determination of
Aromatic in Gasoline
Chemistry - Analysis, Spectrochemical

Sep 48

Quantitative Determination of Aromatic Hydrocarbons C₆-C₉ in Cracking Gasoline by the Method of Combined Dispersion of Light, M. D. Tilichayev, P. A. Akirhin, Ye. G. Treshchova, V. M. Tatevskiy, Cen Inst of Aviation Fuels and Lubricants and Moscow State U, 8 $\frac{1}{2}$ pp

Lavod Lab^o Vol XIV, No 9

Measures frequency and intensities of lines in combination dispersion spectra of aromatics-ethylbenzene, o-, m-, and p-xylenes. Demonstrates possibility of determining these aromatics in cracking gasolines in the presence of alkenes by this method for 60 artificial mixtures. Concentration of individual compounds varied 2-40%.

A 16/49T26

TATEVSKIY, V. M.

CA: 43-2387/h

YUR'YEV, YU. K., TATEVSKIY, V. M. and GRAGEROV, I. I.
J. Phys. Chem (USSR) 22, 783-6 (1948)
Raman spectra of thiophane and its homologs.

PROCESSES AND PROPERTIES ACCT

10

CA

Investigation of the rotation isomerism of 1,2-dibromoethane by the method of Raman spectra. O. J. Aramov, V. P. Zolotarev, and A. V. Kabanov (Moscow State Univ.). *Doklady Akad. Nauk S.S.S.R.* 60, 387-90 (1948).—The equil. const. between the 2 forms of 1,2-C₂H₄Br₂ was detd. from the ratio of the intensities *I* of 2 Raman lines, by $K = A(I_1/I_2) = AK'$, the subscripts 1 and 2 referring, resp., to the frequencies 735 and 659 cm.⁻¹, the 1st belonging to the isomer persisting on freezing, the 2nd to that disappearing on freezing (gauche according to Mitajinska); between -31 and +95° (the variation of *A* in that temp. interval being only about 0.2%, and disregarded), $K = \text{const.} = 1.12 \pm 0.09$, i.e., the ratio of the isomers does not vary in this range. Thus, disappearance of the 2nd isomer can occur only in the solid state or else very close to freezing. In the case of 1,2-C₂H₄ClBr, the ratio of the intensities of the 725 and 667 cm.⁻¹ lines

(the former pertaining to the trans form) remains const. = 1.04 ± 0.09 between -13.6 and +93°, and so does the ratio 725/629 = 2.31 ± 0.14 ; only the ratio 725/696 increases slowly with the temp., from 1.53 to 2.24 between the temp. limits stated. For 1,2-C₂H₄Br, the ratios of the intensities 659/551, at 10, 60, 97, and 128°, are 9.51, 4.43, 3.79, and 3.10. Values of the heat *Q* of transition between the isomers vary in some cases with the Raman lines used: 1,2-C₂H₄Cl (from 735/653) $Q = 140$ cal./mole; 1,2-C₂H₄ClBr (725/667) $Q = 176$, (725/699) $Q = 176$, (725/690) $Q = 160$; 1,2-C₂H₄Br, (659/551) $-2160 = 180$ (between 10 and 128°), (659/551) $-2220 = 230$ (between 10 and 97°), (659/551) $-2680 = 380$ (between 10 and 60°), av. $-2420 = 250$. N. Thon

METALLURGICAL LITERATURE CLASSIFICATION

SECTION DIVISION

MAY 1951

TATEVSKIĬ, V.M.

Hindered rotation in the 1,2-dihaloethane series. O. L. Aronov, V. M. Tatevskiĭ, and A. Frost (Moscow State Univ.): *Doklady Akad. Nauk S.S.S.R.* 60, 1177-80 (1949); cf. preceding abstr. -- The ratio of the intensities I of the Raman lines $753/653 \text{ cm.}^{-1}$ of 1,2- $\text{C}_2\text{H}_4\text{Cl}_2$ was detd. in various solvents: pure liquid 1.12; in CCl_4 (75.2, 50.3, 24.6 mole % of 1,2- $\text{C}_2\text{H}_4\text{Cl}_2$) 1.37, 1.68, 3.17; in C_6H_6 (78.7, 61.4, 23.0) 1.37, 1.23, 2.03; in cyclohexane (76.2, 50.2, 25.4) 1.21, 1.93, 2.70; in C_7H_8 (78.7, 61.4, 23.0) 1.21, 1.47, 2.10; in C_8H_{10} (88.7, 63.4, 25.4) 1.14, 1.13, 1.25; in EtOH (83.0, 51.6) 1.04, 1.20; in PhNO_2 (43.5, 23.1) 0.93, 0.84. For 1,2- $\text{C}_2\text{H}_4\text{Br}_2$, lines 659/551 (the variations at 100 mole % are due to insufficient constancy of the temp.): in CCl_4 (100, 81.0, 43.3) 19.2, 7.0, 7.2, 6.0, 6.2; in C_6H_6 (100, 69.4, 22.6) 7.0, 4.5, 2.7; in cyclohexane (100, 82.5, 51.1, 21.8) 9.1, 8.9, 6.3, 5.3; in C_7H_8 (100, 42.4, 21.4) 7.0, 7.2, 7.3; in C_8H_{10} (100, 46.7, 15.2) 7.0, 6.7, 2.9; in EtOH (100, 69.1, 12.7) 9.1, 8.9, 4.6; in PhNO_2 (100, 41.7, 15.1) 9.1, 4.1, 2.1. For 1,2- $\text{C}_2\text{H}_4\text{Cl}_2$, the dependence of the equil. const. of the 2 rotation isomers ($K = I_1/I_2$) on the solvent is great-

est in the case of nonpolar solvents. Almost no change is found in solvents with a dipole moment close to that of $\text{C}_2\text{H}_4\text{Cl}_2$ itself, e.g. EtOH. Some lowering of K , as compared with the pure liquid, is found in solvents with a dipole moment higher than that of $\text{C}_2\text{H}_4\text{Cl}_2$. The effect of a nonpolar solvent being primarily one of diln., it follows that the 1st form is little polar or nonpolar, while the 2nd form has a higher dipole moment. A solvent with a high dipole moment shifts the equil. in favor of the 2nd form. Strikingly, C_6H_6 behaves more like a solvent of medium polarity, rather than as a nonpolar liquid; this may be

due to its polarizability, giving rise to induced dipole moments. The foregoing considerations are not applicable to 1,2- $\text{C}_2\text{H}_4\text{Br}_2$.
N. Thea

PA 70114

TATEVSKIY, V. M.

Jun 1948

USSR/Chemistry - Ethane, 1,2-Dichloro
Chemistry - Ethane, 1,2-Dibromo

"Rotational Isomers of 1,2-Dichloroethane and 1,2-Dibromoethane in Solution at Different Temperatures,"
O. L. Aronov, V. M. Tatevskiy, A. V. Frost, Moscow
State U imeni M. V. Lomonosov, 3 pp

"Dok Ak Nauk SSSR" Vol LX, No 8

Study the spectra of subject substances to explain the effect of various factors on the equilibrium of rotational isomers in these compounds. Submitted by Acad A. A. Balandin 13 Apr 1948.

70114

USSR/Chemistry - Hydrocarbons, Aromatic Aug 48
Chemistry - Hydrocarbons, Raman Spectra

"Spectra of the Combined Dispersion of Aromatic C₈
Hydrocarbons," Ye. G. Treshchova, V. M. Tatevskiy,
Moscow State U imeni M. V. Lomonosov, 4 pp

"Dok Ak Nauk SSSR" Vol LXI, No 5

Raman spectra of following hydrocarbons with the pu-
rity indicated are reported: ethylbenzene 99.4%,
o-xylene 98.4%, m-xylene (purity undetermined) and
p-xylene 97.2%.

24/49T17

117 AND 3 P. CODERS
PROCESSES AND PROPERTIES INDEX

CA

Spectra of combination scattering of alkenes with an isolated or conjugated system of double bonds V. M. Tatevskii, R. G. Ireshcheva, V. R. Skvarchenko, and R. Ya. Levina. *Zhur. Fiz. Khim.* 35, 657 (1949); *cf. C.A.* 43, 2907. The intensity I of 2 main Raman lines between 1600 and 1700 cm^{-1} is for conjugated double bonds 5-10 times that of nonconjugated. The increase in I is greater the greater the evaluation of the mol. refraction. MR (cf. Shoygin, *ibid.* 42, 2825a) 1,5-Hexadiene: bn 60.5°, d_1^0 0.6015, wp 1.4010, MR 28.00; main Raman lines and, in parenthesis, their I with the assumption that the I of the C=C line of 1-alkenes at 1642 cm^{-1} is 120 per mol. and cm^2 are 1270 (50), 1200 (125), 1410 (85), 1640 (230), 2004 (110), 2015-2001 (120), 2020 (65), 2003 (160), and 2081 (75). 2,5-Dimethyl-1,5-hexadiene: bn 114°, d_1^0 0.7505, wp 1.4363, MR 37.02; 1415 (85), 1434-1442 (90), 1640 (210), 1672 (22), 2001 (250), 2015-2030 (200), and 2046 (170). 2,4-Hexadiene: bn 82°, d_1^0 0.7190, wp 1.4516, MR 30.77; 434 (100), 1150 (150), 1170 (185), 1241 (130), 1270 (80), 1302 (160), 1382 (115), 1440 (135), 1650 (140), 1608 (80), 2004 (100), 2000 (200), 2010 (170), 2008 (115), and 2005 (125). 2,6-Dimethyl-2,4-hexadiene: bn 131-134.5°, d_1^0 0.7636, wp 1.4700, MR 40.83; 205 (60), 307 (60), 840 (80), 863 (85), 1070 (80), 1000 (60), 1150 (75), 1220 (320), 1205 (340), 1382 (60), 1442 (240), 1460 (200), 1650 (130), 1605 (220), 2004 (140), 2015-2000 (450), 2008 (100), and 2081 (150). Line 1302 cm^{-1} belongs to normal alkenes, RCI-CIII. I. I. B.

439.514 METALLURGICAL LITERATURE CLASSIFICATION

13000 131000 132000 133000 134000 135000 136000 137000 138000 139000 140000 141000 142000 143000 144000 145000 146000 147000 148000 149000 150000 151000 152000 153000 154000 155000 156000 157000 158000 159000 160000 161000 162000 163000 164000 165000 166000 167000 168000 169000 170000 171000 172000 173000 174000 175000 176000 177000 178000 179000 180000 181000 182000 183000 184000 185000 186000 187000 188000 189000 190000 191000 192000 193000 194000 195000 196000 197000 198000 199000 200000 201000 202000 203000 204000 205000 206000 207000 208000 209000 210000 211000 212000 213000 214000 215000 216000 217000 218000 219000 220000 221000 222000 223000 224000 225000 226000 227000 228000 229000 230000 231000 232000 233000 234000 235000 236000 237000 238000 239000 240000 241000 242000 243000 244000 245000 246000 247000 248000 249000 250000 251000 252000 253000 254000 255000 256000 257000 258000 259000 260000 261000 262000 263000 264000 265000 266000 267000 268000 269000 270000 271000 272000 273000 274000 275000 276000 277000 278000 279000 280000 281000 282000 283000 284000 285000 286000 287000 288000 289000 290000 291000 292000 293000 294000 295000 296000 297000 298000 299000 300000 301000 302000 303000 304000 305000 306000 307000 308000 309000 310000 311000 312000 313000 314000 315000 316000 317000 318000 319000 320000 321000 322000 323000 324000 325000 326000 327000 328000 329000 330000 331000 332000 333000 334000 335000 336000 337000 338000 339000 340000 341000 342000 343000 344000 345000 346000 347000 348000 349000 350000 351000 352000 353000 354000 355000 356000 357000 358000 359000 360000 361000 362000 363000 364000 365000 366000 367000 368000 369000 370000 371000 372000 373000 374000 375000 376000 377000 378000 379000 380000 381000 382000 383000 384000 385000 386000 387000 388000 389000 390000 391000 392000 393000 394000 395000 396000 397000 398000 399000 400000 401000 402000 403000 404000 405000 406000 407000 408000 409000 410000 411000 412000 413000 414000 415000 416000 417000 418000 419000 420000 421000 422000 423000 424000 425000 426000 427000 428000 429000 430000 431000 432000 433000 434000 435000 436000 437000 438000 439000 440000 441000 442000 443000 444000 445000 446000 447000 448000 449000 450000 451000 452000 453000 454000 455000 456000 457000 458000 459000 460000 461000 462000 463000 464000 465000 466000 467000 468000 469000 470000 471000 472000 473000 474000 475000 476000 477000 478000 479000 480000 481000 482000 483000 484000 485000 486000 487000 488000 489000 490000 491000 492000 493000 494000 495000 496000 497000 498000 499000 500000 501000 502000 503000 504000 505000 506000 507000 508000 509000 510000 511000 512000 513000 514000 515000 516000 517000 518000 519000 520000 521000 522000 523000 524000 525000 526000 527000 528000 529000 530000 531000 532000 533000 534000 535000 536000 537000 538000 539000 540000 541000 542000 543000 544000 545000 546000 547000 548000 549000 550000 551000 552000 553000 554000 555000 556000 557000 558000 559000 560000 561000 562000 563000 564000 565000 566000 567000 568000 569000 570000 571000 572000 573000 574000 575000 576000 577000 578000 579000 580000 581000 582000 583000 584000 585000 586000 587000 588000 589000 590000 591000 592000 593000 594000 595000 596000 597000 598000 599000 600000 601000 602000 603000 604000 605000 606000 607000 608000 609000 610000 611000 612000 613000 614000 615000 616000 617000 618000 619000 620000 621000 622000 623000 624000 625000 626000 627000 628000 629000 630000 631000 632000 633000 634000 635000 636000 637000 638000 639000 640000 641000 642000 643000 644000 645000 646000 647000 648000 649000 650000 651000 652000 653000 654000 655000 656000 657000 658000 659000 660000 661000 662000 663000 664000 665000 666000 667000 668000 669000 670000 671000 672000 673000 674000 675000 676000 677000 678000 679000 680000 681000 682000 683000 684000 685000 686000 687000 688000 689000 690000 691000 692000 693000 694000 695000 696000 697000 698000 699000 700000 701000 702000 703000 704000 705000 706000 707000 708000 709000 710000 711000 712000 713000 714000 715000 716000 717000 718000 719000 720000 721000 722000 723000 724000 725000 726000 727000 728000 729000 730000 731000 732000 733000 734000 735000 736000 737000 738000 739000 740000 741000 742000 743000 744000 745000 746000 747000 748000 749000 750000 751000 752000 753000 754000 755000 756000 757000 758000 759000 760000 761000 762000 763000 764000 765000 766000 767000 768000 769000 770000 771000 772000 773000 774000 775000 776000 777000 778000 779000 780000 781000 782000 783000 784000 785000 786000 787000 788000 789000 790000 791000 792000 793000 794000 795000 796000 797000 798000 799000 800000 801000 802000 803000 804000 805000 806000 807000 808000 809000 810000 811000 812000 813000 814000 815000 816000 817000 818000 819000 820000 821000 822000 823000 824000 825000 826000 827000 828000 829000 830000 831000 832000 833000 834000 835000 836000 837000 838000 839000 840000 841000 842000 843000 844000 845000 846000 847000 848000 849000 850000 851000 852000 853000 854000 855000 856000 857000 858000 859000 860000 861000 862000 863000 864000 865000 866000 867000 868000 869000 870000 871000 872000 873000 874000 875000 876000 877000 878000 879000 880000 881000 882000 883000 884000 885000 886000 887000 888000 889000 890000 891000 892000 893000 894000 895000 896000 897000 898000 899000 900000 901000 902000 903000 904000 905000 906000 907000 908000 909000 910000 911000 912000 913000 914000 915000 916000 917000 918000 919000 920000 921000 922000 923000 924000 925000 926000 927000 928000 929000 930000 931000 932000 933000 934000 935000 936000 937000 938000 939000 940000 941000 942000 943000 944000 945000 946000 947000 948000 949000 950000 951000 952000 953000 954000 955000 956000 957000 958000 959000 960000 961000 962000 963000 964000 965000 966000 967000 968000 969000 970000 971000 972000 973000 974000 975000 976000 977000 978000 979000 980000 981000 982000 983000 984000 985000 986000 987000 988000 989000 990000 991000 992000 993000 994000 995000 996000 997000 998000 999000 1000000

AKISHIN, P. A., KELLE, V. I., TATEVSKIY, V. M., SILAYEV, A. V.

Biophysics.

One mistaken theory of Professor Kobozev. Vest. Mosk. un. 5 no. 8, 1950.

9. Monthly List of Russian Accessions, Library of Congress, November 1953², Uncl.

CA

2

Problems of the chemical structure of hydrocarbons. I. Laws of geometrical configuration of hydrocarbon molecules. V. M. TALEVSKII (Moscow State Univ.). *Vestnik Moskov. Univ. S. No 10, Ser. Fiz.-Mat. i Estestv. Nauk No. 7, 23-28 (1950)*; cf. C.A. 45, 3793d.—The current-resonance theory, represented by Pauling, which attributes the deviations in C—C distances from the standard values for the single, double, and triple bond, to intermediate bond multiplicities (e.g. percent of double-bond character) is assailed. It is contended that any problems posed by the conjugated double or the aromatic bond are irreconcilable with the original structure theory of A. M. Butlerov. Bond distances are not immutable, and are not solely a function of the multiplicity of the bond, but are primarily detd. by the geometric configuration of the atoms involved, whether tetrahedral, plane, or linear. The tacit assumption that in a single bond the C—C distance must be invariably 1.54 Å, irrespective of the geometry is unfounded. A classification of C—C bonds is proposed which is based on the configurations of the valences around each, i.e. on their "coordination nos." The system is admittedly phenomenological, insofar as it does not concern itself with the problem of the phys. meaning of the differences of the valence configurations. In the Pauling curve of interat. C—C distance as a function of the percentage of double-bond character the points for C₁₁H₈, C₁₁H₆, and graphite are correct, but the point for C₁₁H₄ is wrong. An alternative graph is given in which the C—C distance is plotted as a function of α , the fraction of π -bond in bonds of the type $\sigma + \alpha\pi$. In this plot the points for C₁₁H₈ ($\alpha = 1.0$), C₁₁H₆ ($\alpha = 0.5$),

graphite, and butadiene ($\alpha = 0.0$) lie on a straight line. The justification for taking butadiene instead of ethane is that butadiene has a σ bond between C atoms in the same valence state as in C₁₁H₈, C₁₁H₆, and graphite, whereas it is not legitimate to identify the σ bond in C₁₁H₈ with that same bond in C₁₁H₆, and graphite. The linearity of this plot would permit calcn. of the C—C distance in butadiene from the distances in C₁₁H₈, C₁₁H₆, and graphite. In opposition to the viewpoint which sees the partial double-bond character as a consequence of a resonance of structure, intermediate bond multiplicity is interpreted as the result of an addnl. bond in which the electron of each of the atoms involved takes part also in other bonds; consequently, this σ -bond involves only a fraction of the electrons of each atom of the bond. The type of bond in condensed aromatic rings should, with increasing no. of such rings, gradually approach the structure of the basal layer of graphite, i.e. the bond type $\sigma + 1/3\pi$. From this point of view, there should be some difference in the interat. distances of the 10—1 bond in naphthalene and the 14—10 bond in anthracene; however, no data are available at this time to support it. Structure formulas of aromatic compds. are written with each bond represented by a full and a dotted line. The types of C—C bonds here discussed are expressed by one-electron wave functions, ψ in each group, describing sp^2 , sp^1 , and sp states and corresponding, resp. to a C atom with valences in the tetrahedral configuration (type $\rightarrow C \leftarrow$), a C atom with valences at an angle of 120° (type $\rightarrow C \leftarrow$), or the graphite type $\rightarrow C \leftarrow$, and a C atom with valences at an angle of 180° (acetylene or allene types). In condensed aromatic rings, bonds of a type intermediate between those in benzene and in graphite are probable. N. Thom

TATEVSKII, V. M.

Levina, R. Ya. Skvarchenko, V. R., Treshchova, E. G., and Tatevskii, V. M. - "Synthesis of hydrocarbons. IX. Diene hydrocarbons in the synthesis of olefinic hydrocarbons with a central double bond." (p. 419)

SO: Journal of General Chemistry, (Zhurnal Obshchei Khimii), 1950, Vol. 20, No. 3.

TATEVSKY, V.M.

Synthesis of hydrocarbons. X. Partial reduction of conjugated-diene hydrocarbons by sodium in liquid ammonia. R. Ya. Levina, V. R. Skvarczenko, V. M. Tatevsky, and E. G. Treshchova (J. gen. Chem. USSR, 1950, 20, 684-690 [U.S. transl., 721-726]; cf. A., 1950, II, 175).—Partial reduction of $(\text{CHMe:CH})_2$ by Na in liquid NH_3 affords a mixture of the 1:2- and 1:4-dihydro-compounds, CHMe:CHPr^n (60-65%) and $(:\text{CHEt})_2$ (35-40%), and is thus unsuitable for the prep. of the latter; Raman-spectral analysis (details given and discussed) proves the presence of both cis- (30%) and trans-forms (70%). CHMe:CHPr^n is synthesised, for comparison purposes, from $\text{CHMe:CH}\cdot\text{CH}_2\text{Cl}$ and EtMgBr , a mixture of cis- (20%) and trans-forms (80%) being obtained.

Slow addition of $(\text{CHMe:CH})_2$ (16g.) in Et_2O to Na (36 g.) in liquid NH_3 at -50° to -60° , stirring for 5-6 hr., and keeping overnight afford a product. C_6H_{12} (75%), b.p. $67.5^\circ/761$ mm., d_4^{20} 0.6802, n_D^{20} 1.3962 [freed from starting materials by $(:\text{CH}\cdot\text{CO})_2\text{O}$] which is analysed spectrographically (Raman-spectral lines tabulated).

Reaction of aq. HCl with $(\text{CH}_2:\text{CH})_2$ affords 1-chlorobut-2-ene, b.p. $84^\circ/748$ mm., d_4^{20} 0.9282, n_D^{20} 1.4350, and 3-chlorobut-1-ene, b.p. $64^\circ/748$ mm., d_4^{20} 0.8990, n_D^{20} 1.4150; both isomers react with EtMgBr in Et_2O in the usual manner and give identical products, viz., mixed cis- and trans-hex-2-ene. b.p. $67.5 - 67.7^\circ/748.5$ mm., d_4^{20} 0.6805, n_D^{20} 1.3940.

E. S. Stern.

TATEVSKII, V. M.

R. Ya. Levina, V. R. Skvarchenko, E. A. Viktorova, V. M. Tatevskii, and E. G. Treshchova, "Synthesis of hydrocarbons. XI. Partial reduction of diene hydrocarbons with a conjugated system of double bonds by sodium in liquid ammonia." (p. 690)

SO: Journal of General Chemistry, (Zhurnal Obshchei Khimii) 1950, Vol 20, No. 4.

CA

Calculation of the elasticity coefficients of the carbon tetrachloride molecule. A. V. Iogansen and V. M. Tatevskii (Univ. Moscow). *Zhur. Fiz. Khim.* 24, 304-402 (1950).—Splitting of Raman lines of CCl_4 , due to the presence of Cl isotopes is calcd. by Plyashevich's method (*C.A.* 26, 60827) with consts. differing from Stepanov's (*C.A.* 40, 154). The results are in good agreement with the expts. of Langseth (*C.A.* 26, 1189), although Rank and Van Horn (*C.A.* 40, 6994) could not reproduce some of L.'s results. J. J. Bikerman

CA

2

The resonance theory. V. M. Tatevskii (Univ. Moscow). *Zhur. Fiz. Khim.* 24, 597-640(1950).—The theory of chem. resonance is criticized (Pauling *The Nature of the Chemical Bond and the Structure of Molecules and Crystals*, 1944 (cf. C.A. 34, 5734¹); Syrkin and Dyatkina, *The Chemical Bond and the Structure of Molecules*, 1946 (C.A. 41, 8010¹)). J. I. Hikerman

CA

Chemical structure of hydrocarbons and regularities of their geometric configuration. V. M. Tatevskii (M. V. Lomonosov State Univ., Moscow). *Doklady Akad. Nauk S.S.S.R.* 74, 287 (1950). C-C bonds are classified into 12 basic types, characterized by degree of unsat. and valence state of the C atoms. The classes are: ordinary C-C link with tetrahedral C atom and one with planar (120°) valence distribution (alkylethylenes, alkylbenzenes); a link between tetrahedral C and one having linear valence distribution (alkylacetylenes); double link with planar structure and 120° valence angle; single link between 2 planar C atoms with 120° valence angle (conjugated dienes, etc.); triple link with linear valence distribution; link between planar and linear valence forms (conjugated alkynes); single link between 2 atoms with linear and linear (diacetylenes); double link between planar and linear forms (allenes); benzenoid C-C links; graphite C-C links; polynuclear C-C links intermediate between the above 2 forms. G. M. Kozolapoff

1957

CA

Chemical structure of hydrocarbons and regularities in heats of formation. V. M. Tatevskii, V. V. Krutov, and N. A. Mendel'shteyn (M. V. Lomonosov State Univ., Moscow). *Doklady Akad. Nauk S.S.S.R.* 76, 743-4 (1961).
 --The lack of additivity of molar heats of formation in dienes, aromatic compounds, and generally in substances with conjugated unsat. bonds is caused by neglect of the valence states of the C atoms comprising the various C-C links in these mole. When bond energies, used as the basis for mol. calcs., are used for the various valence conditions possible, the results are in excellent agreement with exptl. values.

The basic bond energy units are as follows: H-C 87.717

kcal.; >C-C< 80.488; >C=C< 98.808 (as in C₆H₆);

>C:C< 95.787 (as in allene); >C-C< 63.146; >C-C<

63.304 (as in toluene); >C-C< 99.136 (as in butadiene);

>C-C< 98.174 (as in styrene); >C-C< 87.182 (as in

biphenyl); C-C 87.001 (as in C₆H₆); C=C 84.447

(1-10 link in C₆H₆); C=C 88.301 (C-C in graphite).

The CHI values is calcd. for the alkane series and is assumed const. for all other types, admittedly introducing an error into individual bond energies, but not in the solution of the additivity problem. As a typical result, o-MeEtC₆H₄ gives exptl. ΔH_{298.15} -1761.796 kcal., whereas the calcd. value is -1762.364; for chrysene the values are -2839.92 and -2837.808. rom. G. M. Kosolapoff

1951

TATAEVSKIY V. M.,

FA 172T11

USSR/Chemistry - Thermodynamics

21 Dec 50

"Chemical Constitution of Hydrocarbons and Regularities in the Heats of Formation," V. M. Tataevskiy, Moscow State U imeni M. V. Lomonosov

"Dok Ak Nauk SSSR" Vol LXXV, No 6, pp 819-822

Postulates 10 subtypes of bond CC in alkanes and 3 subtypes (primary, secondary, tertiary) of bond CH having different heat contents. Calcn of heats of formation on basis of principle of additivity then leads to satisfactory results. Same principle of additivity also ought to apply to other values like mol vol and mol refraction, but this must be checked.

172T11

LEVINA, R. YA., FAYNZIL'BERG, A. A., TATEVSKY, V. M., TRESHCHOVA, YE. G.

Carbinols

"Contact dehydration of dimethylallyl carbinol." Uch. zap. Mosk. un. No. 132, 1950.

Monthly List of Russian Accessions, Library of Congress, October 1952. UNCLASSIFIED.

TATEVSKIY, V. M.

"Some Questions of the Theory of Molecular Systems." Sub 28 Dec 51,
Moscow Order of Lenin State U imeni M. V. Lomonosov.

Dissertations presented for science and engineering degrees in
Moscow during 1951.

SO: Sum. No. 480, 9 May 55

TATEVSKI^x, V.M.

Author: Tatevskii, V.M.

Title: Spectroscopy, Ed. by A.V. Frost
189 pp., illus.

Date: 1951, Moscow

Subject: Spectrum Analysis

Available: Library of Congress, Call No: QD55.M3

Sources: Lib. of Cong. Subj. Cat., Apr. June 1952

2

CA

The additive scheme of the heats of formation of hydrocarbons and the problem of the heat of sublimation of graphite. V. M. Tatarskiĭ, R. A. Mendizeritskiĭ, and V. V. Korobov. *Vysokiiĭ Atmosf. Fiz.* 6, No. 5, Ser. Fiz.

Mol. i Kinet. Nash No. 3, 83-6(1951).—The validity of the additive scheme (cf. C.A. 45, 2332b) is independent of the yet uncertain numerical value of the heat of sublimation $\Delta H_{\text{sub}}^{\circ}$ of graphite. A change of that datum can cause only a change of the numerical values of the heats of the different types of C—C bonds and of the C—H bond, but cannot affect the validity of the scheme itself. For this statement to be valid, it is sufficient that, for each class of hydrocarbons, the no. n of C atoms be representable as a linear combination of the nos. n_1, n_2, \dots of C—C bonds of each type, and of m , of the form $n = \alpha n_1 + \beta n_2 + \dots + \gamma m$, where the coeffs. $\alpha, \beta, \dots, \gamma$ are consts. Fulfillment of that condition is illustrated by the examples of alkanes, cycloalkanes, alkenes, and alkynes. N. Thon

CA

3

Raman spectra of α -alkenes. O. D. Ul'yanova and V. M. Yalovskii, *Vestnik Mosk. Univ.* 6, No. 6, Ser. Fiz.-Mat. i Estet. Nauk No. 3, 87-9 (1951).—The following Raman frequencies, in cm^{-1} (intensities on a 10 scale), were observed in 5 hrs. exposure in excitation with 4358 Å., dispersion 13 Å./mm. 1-C₆H₁₀, 242(2), 263(1), 315(0), 358(0), 400(0), 478(0), 638(2), 635(0), 746(0), 810(3), 875(1), 895(2), 912(5), 931(4), 984(2), 1056(2), 1108(3), 1216(3), 1290(8), 1413(7), 1441(7), 1642(10), 2728(2), 2849(3), 2883(6), 2910(3), 2935(5), 2962(6), 2990(7); 1-C₇H₁₂, 243(1), 281(0), 339(0), 388(0), 435(0), 490(0), 558(1), 635(1), 679(0), 733(0), 782(3), 833(0), 918(4), 925(3), 990(2), 1069(3), 1114(3), 1214(2), 1292(8), 1413(8), 1452(7), 1641(10), 2843(3), 2888(7), 2927(3), 2953(4), 2992(7), 3400(4); 1-C₈H₁₄, 221(0), 246(1), 404(0), 438(0), 469(1), 635(1), 711(1), 728(0), 794(1), 830(0), 834(0), 858(1), 885(3), 906(4), 938(0), 967(1), 998(2), 1028(2), 1061(3), 1081(2), 1111(3), 1210(2), 1298(8), 1372(2), 1416(7), 1439(8), 1469(8), 1641.7(10), 2888(7), 2928(2), 2954(4), 2998(7), 3060(5). Abs. intensities per 1 mole, $I_{0.1}$, were detd. from the measured intensities I_1 of the alkene, by comparison with the measured intensities I_2 of cyclohexane added to alkene, by $I_{0.1} n_1 / I_{0.2} n_2 = I_1 / I_2$, where n_1 and n_2 = mol. concns. of the alkene and of cyclohexane, resp., and the abs. $I_{0.1}$ for the line 802 cm^{-1} of cyclohexane was taken = 250 units/mole. Values of $I_{0.1}$ for the strongest lines of 1-C₆H₁₀, 1-C₇H₁₂, and 1-C₈H₁₄, are: 910 cm^{-1} , 16.94, 24.15, 16.96; 1296 cm^{-1} , 26.52, 110.35, 65.30; 1413 cm^{-1} , 21.06, 19.47, 29.70; 1440 cm^{-1} , 21.30, 41.50, 75.05; 1642 cm^{-1} , 111.1, 123.6, 121.2. There are minor differences between these Raman spectra and those observed by Bashulin and Sterin (*Izv. AN SSSR, Ser. fis. 11, No. 4, 481 (1947)*). Analytical application of the Raman spectra to the detn. of 1-C₆H₁₀ in mixt. with C₇H₁₂ and cyclohexane was very satisfactory with 35 and 8% 1-C₆H₁₀; the lower limit of the detn. is 3% 1-C₆H₁₀. N. Thon

CA

2

Some aspects of the theory of chemical structure proposed
by A. M. Butlerov. V. M. Tatevskii. *Vestnik Moskov.
Univ.* 6, No. 6, Ser. Fiz.-Mat. i Estestv. Nauk No. 4, 53-70
(1951).—Review with numerous quotations from Butlerov's
publications supporting his priority as the true founder of
structural theory of organic chemistry. G. M. Kosolapoff

Intensity of the band of the valence vibrations of the C=C bond in the Raman spectra of hydrocarbons. P. A. Alshin and M. M. Tatarskiy, Moscow State Univ., Vestnik MGU, Ser. Fiz.-Mat. i Estestv. Nauk No. 1, 1951, No. 1, p. 15. The Raman line intensities were determined against standard CHCl_3 ($\nu_{\text{C-Cl}}$ at 300 cm^{-1}), cyclohexane (800), C_2H_4 (1623) and C_2H_2 ($\nu_{\text{C-C}}$ at 2100 cm^{-1}). The differential intensities are expressed on a scale in which that of the cyclohexane line at 800 cm^{-1} is = 250 units/mole cm^{-1} , and integral intensity relative to that line = differential and integral intensity in cm^{-1} . Data (C=C frequency in cm^{-1} , differential and integral intensity) are: Methylcyclopentane 1676, 125, 120; 1-butene 1642, 129, 129; propylene 1671, 124, 123; 2-methyl-2-butene 1644, 127, 125; 1,3-butadiene 1658, 131, 125; 1,2-cyclohexadiene 1644, 127, 125; 1,2,3-cyclohexatriene 1653, 123, 123; cyclopentadiene 1657, 129, 127; 1-methyl-cyclopentene 1655, 115, 120; 1,3-butadiene 1657, 129, 140; vinylcyclopentane 1657, 129, 140; allylcyclopentane 1676, 122, 141; 1,3-butadiene 1671, 124, 123; 2-ethyl-1-butene 1654, 125, 129; 1-methyl-cyclohexene 1676, 122, 141; 1,3-butadiene 1645, 122, 125; methylcyclohexene 1651, 120, 123; vinylcyclohexane 1641, 113, 125; allylcyclohexane 1641, 134, 126; linoleene 1644, 122, 130; 1,6-heptadiene 1642, 127, 147. The differential intensity appears to be remarkably constant (125-130) units, independently of the structure of the hydrocarbon. Where the C=C band is split into 2 components, as a result of internal rotation or Fermi resonance, the sum of their intensities nearly remains constant. From the differential and the integral intensities, the half width of the C=C line appears constant, $\sim 2 \text{ cm}^{-1}$.

CA

TATYNSKII, V. M.

"Isomerisation of unsaturated hydrocarbons over metallic oxides. XII. Preparation of methyl pentadienes by dehydration of dimethyl allyl acrylate over chromium oxide-on-alumina." by R. Y a. Levina, A. A. Fainzil'berg, V. M. Tatyanski, E. G. Beresina. (p.233)

SO: Journal of General Chemistry (Zhurnal Obshchei Khimii) 1951, Volume 21, No. 2

TATEVSKIY, V. M.

184T36

USSR/Chemistry - Resonance Theory Feb 51

"Letter to the Editor Regarding V. M. Tatevskiy's Article 'On the Theory of Resonance,'" G. V. Chelintsev, Moscow

"Zhur Fiz Khim" Vol XXV, No 2, pp 239, 240

Criticizes V. M. Tatevskiy for (1) usurpation of Chelintsev's priority in criticism of theory of resonance, giving Chelintsev no credit; (2) incomplete criticism of theory based on quantum mech aspects only; (3) acceptance of postulates of dislocation and diffusion of electrons, which in effect retained resonance theory; (4) acceptance
184T36

USSR/Chemistry - Resonance Theory Feb 51
(Contd)

of mesomerism; (5) nonacceptance of Chelintsev's orbital theory.

184T36

184T37

TATEVSKIY, V. M.

USSR/Chemistry - Resonance Theory Feb 51

"Certain Problems of A. M. Butlerov's Theory of Chemical Structure. II. Chemical Structure of Hydrocarbons and Regularity in Energies of Formation," V. M. Tatevskiy, Moscow State U Iment V. M. Lomono-
sov

"Zhur' Fiz Khim" Vol XXV, No 2, pp 241-254

Rejects mol formulas of A. M. Butlerov as too rigid in light of newer sci knowledge. Calls resonance theory idealistic. States methods for calcn of resonance energy are faulty. Concept of additivity agrees with results from calcn of heat of formation

184T37

USSR/Chemistry - Resonance Theory (Contd) Feb 51

of aromatic hydrocarbons by orbital method of Slater's method. Quantum chem calcns show practically complete additivity.

184T37

USSR/Chemistry - Isotopes

Mar 51

"Calculation of the Equilibria of Isotopic Exchange Reactions," V. M. Tatevskiy, Lab Mol Spectroscopy, Moscow State U imeni M. V. Lomonosov

"Zhurnal Fiz Khim" Vol XXV, No 3, pp 261-273

Obtained expression for equil consts (Kp) depending only on numbers of symmetry by proposed new theorem and 2 new variants of statistical method for calcn of isotope equls. Treated temp dependence of Kp and showed approx method for its calcn for certain temps using numbers

185T4

USSR/Chemistry - Isotopes (Contd)

Mar 51

of symmetry, geom forms of mols, and collection of fundamental frequencies for only 1 isotopic mol of each type. Ratio of products of chief moments of inertia of isotopic mols depends only on and can be calcd from mass, geom form. Illustrated reactions include isotope-substituted forms of CCl4, CH4, H2O, H2S.

185T4

185T4

TATEVSKIY, V. M.

TATEVSKIY, M. V.

185T5

USSR/Chemistry - Isotopes

Mar 51

"Use of Lemmata of Similarity for the Calculation of the Vibration Spectrum of Isotopic Molecules," M. V. Tatevskiy, Lab Mol Spectroscopy, Moscow State U imeni M. V. Lomonosov

"Zhur Fiz Khim" Vol XXV, No 3, pp 274-282

Indicates method for use of lemmata of similarity for calcn of vibration frequencies of isotopic mols XY_n using exptl data for mols substituted with other isotopes. Examples include deuterium- and tritium-substituted H_2O , H_2S , H_2Se , NH_3 , PH_3 , AsH_3 , CH_4 , SiH_4 , GeH_4 , NH_4^+ , and WO_4^{2-} , IO_3^- , ReO_4^- , substituted with O17.

185T5

TATEVSKIY, V. M.

USSR/Chemistry - Antiknock Fuels

Oct 51

"Spectra of Combination Scattering of Certain C_6-C_{11} Isoparaffines With Tertiary Carbon Atoms," Ye. G. Treschova, V. M. Tatevskiy, T. I. Tantsyeva, A. A. Paynill'berg, R. Ya. Levina, Moscow State U imeni M. V. Lomonosov.

"Zhur Fiz Khim" Vol XXV, No 10, pp 1239-1247.

Measurements of number and intensity of lines in spectra of combination scattering of 2, 4-dimethylhexane, heptane, octane, and nonane, 2, 4, 5-trimethylhexane, 2, 4, 6-trimethylheptane, and 2, 4, 7-trimethyloctane, in agreement with theoretical conclusions of B. I. Stepanov, showed that different types of branchings of alkane C skeleton are represented by sp characteristics of spectrum.

PA 194T18

TATAYEVSKIY, V. H.

178T15

USSR/Chemistry - Hydrocarbons

1 Feb 51

"Intensity of the Band of the C=C Valency Vibration in Combination Dispersion Spectra of Hydrocarbons," P. A. Akishin, V. M. Tatayevskiy, Moscow State U imeni M. V. Lomonosov

"Dok Ak Nauk SSSR" Vol LXXVI, No 4, pp 527-530

Lists frequencies in cm^{-1} produced by C=C bonds, differential intensities per mol in cm^{-1} , and integral intensities per mol for 21 unsatd hydrocarbons of different constitution. General relationships established on basis of these data can be used in quant mol analysis and for detg structural elements (isolated C=C bonds).

178T15

TATEVSKIY, V.M.

USSR/Chemistry - Fuels

1 May 51

"Chemical Structure of Hydrocarbons and Regularities in the Heats of Formation," V. M. Tatevskiy, V. V. Korobov, E. A. Mendzheritskiy, Moscow State U imeni M. V. Lomonosov

"Dok Ak Nauk SSSR" Vol LXXVIII, No 1, pp 67-69

The authors studied the heats of formation of 44 straight and branched-chain hydrocarbons. They give a table depicting the increments in the heat of formation for the addn of single carbon atoms to the chain. Values for $\Delta H^{\circ}_{C_nH_{2n+2}(at)}$ were calcd on an additive basis for the temps 0 and 298.16° K.

21713

TATEVSKIY, V.

A.V.Frost; obituary. Vest.Mosk.un. 7 no.12:131-137 D '52. (MLRA 7:9)
(Frost, Andrei Vladimirovich, 1906-1952)

TATEVSKIY, V. M.

USSR/Chemistry-Hydrocarbons

Feb 52

"Synthesis of Hydrocarbons. XVII. Homomesitones in the Synthesis of Isoparaffinic Hydrocarbons," R. Ya. Levina, N. P. Shusherina, Ye. G. Treshchova, V. M. Tatevskiy, Lab of Org Chem imeni Acad N. D. Zelinskiy, Moscow State U

"Zhur Obshch Khim" Vol XXII, no 2 pp 199-208

Developed method to synthesize difficulty obtainable isoparaffins 3, 5-dimethylheptane (I), 3-methyl-5-ethylheptane (II), 3-methyl-5-ethyloctane (III), 3-methyl-5-ethylnonane (IV), 2, 3, 4-trimethylhexane (V), 3, 4, 5-trimethylheptane (VI), none described in the literature. Synthesized I-IV by reaction of mixt of "homomesitones" 3-methylheptene-3-one-5 and 3-methylheptene-2-one-5 (products of condensation of methylethylketone with CaC_2) with org Mg compds, then dehydration of products (unsatd tert alcs) and catalytic hydrogenation of resulting diene hydrocarbons. V and VI were synthesized in same way with exception that "homomesitones" in this case were 3, 4-dimethylhexene-3-one-5 and 3, 4-dimethylhexene-2-one-5 (products of condensation of methylethylketone with HCl).

209T14

TATAYVSKIY, V. H.

261T37

USSR/Chemistry - Hydrocarbons

Sep 52

"Raman Spectra of Hydrocarbons of Various Types: III. Raman Spectra of Some Alkyldienes With a System of Isolated or Conjugated Double Bond," Ye.G. Treshchova, V.M. Tatayvskiy, R. Ya. Levina, A.A. Faynzil'beris, and Ye.A. Viktorova, Moscow State U

Zhur Fiz Khim, Vol 26, No 9, pp 1266-1271 5

The Raman spectra of the following alkyldienes were studied: 2-methyl-pentadiene-1,4; 2,3-dimethylbutadiene-1,3; 2-methylpentadiene-2,4; heptadiene-2,4; octadiene-2,4. The intensities of the lines of the

261T37

spectrum were measured photometrically in a single objective scale. In the spectrum of alkyldienes with isolated double bonds there were in the range 1,600-1,680 cm⁻¹ either one strong line with an intensity approx twice as great as that of the alkenes or two strong lines, the total intensity of which was approx twice as great as that of the alkenes. In the spectrum of alkyldienes with conjugated double bonds, the intensity of the characteristic lines in the range of 1600-1680 cm⁻¹ surpasses the intensity of the characteristic line in this field in the case of alkenes. A comparison was made of the spectra of alkyldienes of normal structure with a system of conjugated double bonds: hexadiene-2,4, heptadiene-2,4, and octadiene -2,4. The authors established the approx recurrence of a series of lines, by frequency and intensity, for all three alkyldienes.

261T37

TATEVSKIY, V.M.; FROST, A.V., professor, redaktor.

[Chemical structure of hydrocarbons and the regularity of their
physicochemical properties] Khimicheskoe stroenie uglevodorodov
i zakonmernosti v ikh fiziko-khimicheskikh svoistvakh, [Moskva]
Izd-vo Moskovskogo universiteta, 1953. 319 p. (MLRA 6:12)
(Hydrocarbons)

TATEVSKIY, V.M.

③

Chemical structure of hydrocarbons and the regularities in the heats of combustion of liquid alkanes and alkylbenzenes. V. M. Tatevskii and Yu. A. Pentin. *Vestnik Moskov. Univ. B, Ser. Fiz.-Mat. i Estestven. Nauk.* No. 3, 69-72 (1953); cf. *C.A.* 45, 8869b; 46, 5906i. —Heats of combustion calcd. from the previously developed expression, $-\Delta H = \sum n_i A_i$, agreed well with ΔH 's detd. by expt. for liquid alkanes, $C_n H_{2n+2}$. ΔH calcd. for 2,3-Me₂C₈H₁₈ and 2,2,4-Me₃C₈H₁₈ were higher, and those for 2,2,3,4-Me₄C₈H₁₈ and 2,2,4,4-Me₄C₈H₁₈ were lower, than the exptl. values. The reason for these exceptions could not be explained.

G. Meguerian

10-13-54 I.J.P.

TATEVSKIY, V.M.; PENTIN, Yu.A.

Chemical structure of hydrocarbons and mechanisms of heat of combustion of liquid alkanes and alkylbenzenes. Vest.Mosk.un. 8 no.5:69-72 *Ky* '53.
(MLDA 6:8)

1. Kafedra fizicheskoy khimii. (Hydrocarbons) (Heat of combustion)

TATEVSKIY, V.M.; KARAPET'YANTS, M.Kh. [authors]; TILICHEYEV, M.D. [redaktor];
KIRBYEV, V.A. [reviewer].

"Physicochemical properties of individual hydrocarbons." Edited by M.D.
Tilicheev. Reviewed by V.A.Kireev. Zhur.fiz.khim. 27 no.6:939-940 Je
'53. (MLRA 6:7)
(Hydrocarbons)

TATAYEVSKIY, V. M.

USSR/Chemistry - Hydrocarbons

Oct 53

"Raman Spectra of Hydrocarbons of Different Types. IV.
Raman Spectra of Some Branched Alkanes Containing
Tertiary Carbon Atoms," Ye. G. Treshchova, V. M.
Tatayevskiy, A. A. Faynzil'berg, N. P. Shusherina,
R. Ya. Levina, Moscow State U im M. V. Lomonosov

Zhur Fiz Khim, Vol 27, No 10, pp 1564-71.

Detd the Raman spectra of 2,3-dimethylhexane, 2,3-
dimethylheptane, 2,3-dimethyloctane, 2,3,6-trimethyl-
heptane, 3,5-dimethylheptane, 3-methyl-5-ethyloctane,
3,5-dimethylheptane, and 5-methyl-5-ethyloctane. From

272T15

the data obtained, drew conclusions on the frequencies
which correspond to definite types of branching.

AKISHIN, P.A.; ~~TATYEVSKIY, V.M.~~

Intensity of bands of valence vibration of the $C\equiv C$ bond in Raman spectra
of hydrocarbons. Doklady Akad. Nauk S.S.S.R. 89, 287-9 '53. (MIRA 6:3)
(GA 47 no.20:10349 '53)

1. M.V.Lomonosov State Univ., Moscow.

TATEVSKIY, V.M.

USSR:

539.132

3714. Rotational isomerism of 1,2-dichloroethane and 1,2-dibromoethane. YU. A. PERMIN, YA. S. BOBICH, D. B. GUREVICH AND V. M. TATEVSKIY. Dokl. Akad. Nauk SSSR, 89, No. 3, 435-8 (1953) In Russian. English translation, U.S. National Sci. Found. NSF-47-38.

The Raman spectra have been recorded photoelectrically at various temperatures between 12 and 117°C. From the relative intensities of the two peaks in each spectrum corresponding to the C-halogen bond vibrations of the two isomers of each compound (at 654 and 754 cm⁻¹ for dichloroethane, and

at 551 and 659 cm⁻¹ for dibromoethane) at the different temperatures, a value close to 9 cal/mole was derived for the heat of isomerization of the dichloro, and of 770 ± 60 cal/mole for the dibromo derivative; this latter figure agrees well with some other published figures, and it is concluded that the value of 2420 cal/mole, published by Prox (Dokl. Akad. Nauk SSSR, 60, 387 (1948)) is erroneous.

R. C. MURRAY

9
Cher

RCW
RM

TATEVSKIY, V.M.

The Committee of Stalin Prizes (of the Council of Ministers USSR) in the fields of science and inventions announces that the following scientific works, popular scientific books, and textbooks have been submitted for competition for Stalin Prizes for the years 1952 and 1953. (Sovetskaya Kultura, Moscow, No. 22-43, 20 Feb - 3 Apr 1954)

| <u>Name</u> | <u>Title of Work</u> | <u>Nominated by</u> |
|-----------------|---|--|
| Tatevskiy, V.M. | "Chemical Structure of Hydrocarbons and the Laws of their Physicochemical Properties" | Moscow State University imeni M.V. Lomonosov |

SC 4-3064, 7 July 1956

TATEVSKIY, V.M.
USSR/Chemistry

FD-774

Card 1/1 : Pub 129 11/24

Author : Tatevskiy, V. M.; Pentin, Yu. A.

Title : ~~Chemical structure and regularities in the heats of combustion of liquid alkanes and alkyl benzenes~~
: Chemical structure and regularities in the heats of combustion of liquid alkanes and alkyl benzenes

Periodical : Vest. Mosk. un., Ser. fizikomat. i yest. nauk, Vol 9, No 2, 81-84, Mar 1954

Abstract : Listed the experimental and calculated values for the heats of combustion of liquid alkanes and of alkyl benzenes. Three tables, Four references (all USSR).

Institution : Chair of Physical Chemistry

Submitted : September 17, 1953

USSR/Chemistry - Physical

TATEVSKIY, V.M.

FD-1677

Card 1/1 : Pub. 129-2/25

Author : Tatevskiy, V. M.; Pentin, Yu. A.

Title : Chemical structure and physicochemical properties of molecules

Periodical : Vest. Mosk. un Ser. fizikom. i vest. nauk, Vol 10, 21-32, Feb 1955

Abstract : The physicochemical properties of a molecule are related to its structure. Molecules are arranged into classes which are determined by the position in the periodic table, bond angle, and other factors that affect the interatomic distances. The classes are further subdivided into types and the types into subtypes. Thus, hydrocarbon molecules were arranged systematically by structure and correlated with physicochemical properties such as molecular volume, refraction, heat of formation, etc. Ultimately, if the physicochemical properties and the proper constants are known, the structure of a molecule can be predicted. Large table; graphs. No references.

Institution : Chair of Physical Chemistry

Submitted : September 17, 1954

FD-2169

USSR/Chemistry - Physical

Card 1/1 Pub 129-9/20

Author : Pentin, Yu A., and Tatevskiy, V. M.

Title : Infrared absorption spectra of certain alkyl halides and the question of rotational isomerism

Periodical : Vest. Mos. un., Ser. fizikomat. i yest, nauk, 10, No 2, 63-72, Mar 1955

Abstract : Obtained infrared absorption spectra in the region $400-1500\text{ cm}^{-1}$ for propyl chloride, propyl bromide, isobutyl bromide, n-butyl bromide, isoamyl bromide, 1,2-dichloroethane, 1,2-dibromoethane, 1,2-dibromopropane, and 1,2-disbromobutane. Selected lines for investigating the effects of temperature on relative line intensity and the determination of the relative stability of the rotational isomers of the above compounds. Graphs, formulas. Sixteen references (six USSR; fifteen since 1940)

Institution : Chair of Physical Chemistry

Submitted : October 20, 1954

USSR/Chemistry - Physical

FD-2170

Card 1/1 Pub 129-10/20

Author : Pentin, Yu. A., and Tatevskiy, V. M.

Title : Energy differences and possible configuration of rotational isomers of certain alkyl halides

Periodical : Vest. Mosk. un., Ser. fizikommat. 1 yest, nauk, 10, No 2, 73-82, Mar 1955

Abstract : Studied the effect of temperature (24-80 degrees) on the optical density of infrared absorption bands and calculated the energy differences of rotational isomers of gaseous propyl chloride and bromide, n-butyl and isoamyl bromides, 1,2-dichloro and 1,2-dibromoethane, and 1,2-dibromopropane. Also conjectured on the possible configurations of the rotational isomers of the above compounds and drew conclusions on their relative stability from determined factors. Tables, graphs, diagrams. Twelve references (six USSR; all since 1940).

Institution : Chair of Physical Chemistry

Submitted : October 20, 1954

TATEVSKIY, V.M.

USSR/Chemistry - Physical chemistry

Card 1/1 Pub. 22 - 31/49

Authors : Tatevskiy, V. M.

Title : Valent carbons in molecules. A concept on the valent state of atoms

Periodical : Dok. AN SSSR 101/3. 515-516, Mar 21, 1955

Abstract : Data are presented regarding valent carbons of various molecules and the valent state of atoms. It is proven that the geometrical configuration of atoms does not experience any basic changes during the substitution of the given central atom by an atom of another element but, of the same group. The configuration remains unchanged (qualitatively) even during the change of one or several atoms directly bound with the given central atom. It is explained that the geometrical configuration depends, basically, upon the state of the central atom (its valence, series and multiplicity of chem. bonds). Table.

Institution :

Presented by : Academician A. N. Frumkin, October 19, 1954

USSR/Chemistry - Physical chemistry

Card 1/2 Pub. 22 - 31/51

Authors : Tatevskly, V. M.

Title : ~~Types of chemical bonds of atoms and interatomic spaces in molecules of halogen derivatives~~
Types of chemical bonds of atoms and interatomic spaces in molecules of halogen derivatives

Periodical : Dok. AN SSSR 101/5, 901-902, Apr 11, 1955

Abstract : The three trivalent states in which a tetravalent carbon atom in molecules of organic compounds can exist are described. It is shown that all chemical carbon-halide bonds in halogen derivatives can be divided into three basic types depending of course upon the valent state of the carbon atom. The interatomic spaces were found to remain approximately constant for bonds belonging to one of the basic types; but, they are different for bonds of different types. Each bond type

Institution :

Presented by : Academician N. N. Semenov, October 19, 1954

Card 2/2

Pub. 22 - 31/51

Periodical : Dok. AN SSSR 101/5, 901-902, Apr 11, 1955

Abstract : was found to be characterized by a specific interatomic space. The effect of atoms which do not directly participate in the formation of the given chemical bond is explained. Five references: 3 USSR and 2 USA (1934-1953). Table; graphs.

VM

As applied to the theory of free energy. The discussion is
concerned with the stability of various substances, in
particular of various mixtures of the elements H, C, O, N,
S, Pb, N, O, As and S. Mol. spectra of org. componds. are
interpreted in terms of free energy of formation of 39 alkanes
at 0°, 298.16°, and 600°K., and of 30 alkanes and 11 alkyl-
cyclohexanes at 298.16 and 600°K. The exptl. and com-
puted values are related to the equations for small

TATEVSKIY, V.M.

*Platz
1
Bauk
W*

Constructional changes in the registration system of the spectrograph DFC-4. A. A. Mal'tsev, M. E. Misyurevich, V. A. Koryazhkin, and V. M. Tatevskii (State Univ., Moscow). *Prilozheniya k Zhurn. Eksperimenty 1970, No. 2*, 124-3. The authors describe the construction of a new registration system for the spectrograph DFC-4. The modification of an automatic registration system increases the resolving power of the instrument. Werner Jacobsen

Г. А. ТЕУСКВИЧ

Раман спектры конденсированных систем

h

TATEVSKI, V.M.

SHAKHPARONOV, M.I.; TATEVSKIY, V.M.

Concerning the discussion on the theory of molecular structure.
Zhur.fiz.khim. 30 no.9:2122-2123 S '56. (MIRA 9:12)

1. Moskovskiy gosudarstvennyy universitet imeni M.V.Lomonosova.
(Molecules)

TATEVSKIY, V. M.

USSR/Physical Chemistry - Molecules. Chemical Bonds.

B-4

Abs Jour: Ref Zhur-Khimiya, No 5, 1957, 14434

Author : Yu. A. Pentin, V. M. Tatevskiy

Inst : -

Title : Energy differences of inverted isomers of certain halogeno derivative hydrocarbons

Orig Pub: Dokl. AN SSSR, 1956, 108, No 2, 290-293

Abstract: With the aim of making a study of inverted isomery in alkyl halides 1,2-dihalogeno derivative hydrocarbons, the following spectra were examined: infra-red, comb. of solutions propylchloride (I), propyl bromide (II), n-butyl bromide (III), isoalyl bromide (IV), 1,2-dichloroethane (V), 1,2-dibromoethane (VI), 1,2-dibromopropane (VII). Infra-red spectra (in the region of 400-1500 cm^{-1}) were examined in solid, liquid and gaseous states, and the comb. of solutions spectra - in the liquid phase at various temperatures. The relation of frequencies to the trans- or turned inverted isomers was checked by an examination of the

Card 1/4

USSR/Physical Chemistry - Molecules. Chemical Bonds.

B-4

Abs Jour: Ref Zhur-Khimiya, No 5, 1957, 14434

Abstract: influence of crystallization on the spectra (during crystallization, lines corresponding to the turned isomer should appear). Energy differences ΔU between the turned and trans-isomers, which are determined by temperature progress of special line intensities, are equal to (in cal/mole, first number corresponds to the gaseous, second - to the liquid state): I -50 ± 70 ; -300 ± 150 ; II -280 ± 100 , -440 ± 100 ; III -50 ± 150 , -100 ± 100 ; IV -490 ± 80 ; $+20 \pm 100$; V 1240 ± 50 , 0 ± 60 ; VI 1780 ± 50 , 770 ± 60 . VII contains 3 different inverted isomers, in which the atom Br of the group CH_2Br is found in the trans-position in relation corresponding to atom H(I), group $\text{CH}_3(1a)$ and atom Br(2) of the group $\text{C}(\text{CH}_3)\text{HBr}$; $u_{1a} - u_2 = \sim 2000$, (?) 1020 ± 250 , $u_1 - u_2 = 1330 \pm 120$, 330 ± 200 ; $u_1 - u_{1a} = -300 - 500$ (?), -670 ± 100 (the authors assume that $u_2 < u_1 < u_{1a}$).

Values of ΔU were determined as the mean values from measurements, conducted for different pairs of bands and

Card 2/4

USSR/Physical Chemistry - Molecules. Chemical Bonds.

Abs Jour: Ref Zhur-Khimiya, No 5, 1957, 14434

B-4

Abstract: the distribution of the C atoms removed from the halogen, so that the isomers decompose into 2 groups, in each of which the C-Hal frequency is practically identical. It was shown, that the experimentally determined value ΔU represents the arithmetical mean value of the energy differences of isomers which belong to these two groups. An equation is also obtained which ties in the observed relation of intensities with the energy differences of all isomers of both groups, which thus may be determined in principle if the intensity relationships are measured in a sufficiently wide interval of temperatures.

Card 4/4

Category: USSR / Physical Chemistry
Thermodynamics, Thermochemistry. Equilibrium. Physico-chemical analysis. Phase transitions.

B-8

Abs Jour: Referat Zhur-Khimiya, No 9, 1957, 29896

Author : Tatevskiy V. M.

Inst : Academy of Sciences USSR

Title : Heat of Evaporation, Vapor Tension and Boiling Point of Paraffinic Hydrocarbons (Alkanes)

Orig Pub: Dokl. AN SSSR, 1956, 109, No 1, 131-134

Abstract: To calculate heat of evaporation, vapor pressure and boiling point of alkanes, the values of λ and b in the equation $\lg p = - \lambda/T + b$ (1) (λ -- molal heat of evaporation, not corrected for non-ideal nature of gas), are calculated analogously to the previously described manner of calculating the other physico-chemical quantities of alkanes (RZhKhim, 1956, 167), in accordance with the formulas $\lambda = \sum n_i \lambda_i$ (2) and $b = \sum n_i b_i$ (3). Values of λ_i and b_i are calculated, on the basis of available experimental data

Card : 1/2

-18-

Category: USSR / Physical Chemistry
Thermodynamics, Thermochemistry. Equilibrium. Physico-
chemical analysis. Phase transitions.

B-8

Abs Jour: Referat Zhur-Khimiya, No 9, 1957, 29896

for various temperatures within the range of 40-150°; for the same temperatures are calculated, and compared with the experimental values, the λ and b of 31 alkanes. Boiling points of 65 alkanes, at p = 760 mm Hg, calculated in accordance with formulas 1, 2 and 3, deviate, in most cases, from the experimental data by less than 1°, and only in one instance the difference amounts to 3°.

Card : 2/2

-19-

TATEVSKIY, V..M.

"Methods for the Statistical Computation of the Isotopic Exchange Equilibrium in the Gaseous Phase."

Problems Kinetics and Catalysis, 7. 9, Isotopes in Catalysis, Moscow, Izd-ro AN SSSR, 1957, 142p.

Most of the papers in this collection were presented at the Conf. on Isotopes in Catalysis which took place in Moscow, Mar 31- Apr 5, 1956.

TATEVSKIY, V. V.

PRIKHOT'KO, A. F.

24(7)

3

PHASE I BOOK EXPLOITATION SOV/1365

L'vov. Universitet

Materialy I Vsesoyuznogo soveshchaniya po spektroskopii. t. 1: Molekulyarnaya spektroskopiya (Papers of the 10th All-Union Conference on Spectroscopy. Vol. 1: Molecular Spectroscopy) [L'vov] Izd-vo L'vovskogo univ-ta, 1957. 499 p. 4,000 copies printed. (Series: Its: Fizichnyy zbirnyk, vpp. 3/8/)

Additional Sponsoring Agency: Akademiya nauk SSSR. Komissiya po spektroskopii. Ed.: Jaser, S.L.; Tech. Ed.: Saranyuk, T.V.; Editorial Board: Lavstern, G.S., Academician (Resp. Ed., Deceased), Neporent, B.S., Doctor of Physical and Mathematical Sciences, Pabelinskiy, I.L., Doctor of Physical and Mathematical Sciences, Fabrikant, V.A., Doctor of Physical and Mathematical Sciences, Korotkiy, V.G., Candidate of Technical Sciences, Rayskiy, S.M., Candidate of Physical and Mathematical Sciences, Klimovskiy, L.K., Candidate of Physical and Mathematical Sciences, Miliyanchuk, V.S., A. Ye., Candidate of Physical and Mathematical Sciences, and Glauberman,

Card 1/30

Pektin, Yu. A., V.M. Tatevskiy, and B.A. Pozdyshev. Study of Rotational Isomerism by Means of Spectroscopy

300

Movak, I.I., and Ya. S. Solov'yev. Rotational Isomerism and the Effect of Temperature on the Infrared Absorption Spectra of Some Paraffins

419

Postovakaya, A.F., I.A. Salimov, A.S. Kur'minskiy, and V.M. Tatevskiy. Variation in Structure of Sodium Butadiene Number in the Process of Light Oxidation

423

Klauzen, N.A., and B.A. Dogadkin. Infrared Spectroscopy Used to Study the Interaction of Rubber and Sulfur

426

Klovokhotova, N.A. Study of the Chemical Variations of Tetrafluoroethylene ("teflon") Under the Influence of Ionizing Radiation by Means of Infrared Spectroscopy

430

Mel'son, K.V., and I. Ya. Podubnyy. Spectroscopic Study of the Microstructure of Some Diene Polymers

433

Card 27/30

TATEVSKIY, V.M.

PRIKHOT'KO, A.F.

24(7) 3 PHASE I BOOK EXPLOITATION 807/1365

L'vov. Universytet

Materialy I Vsesoyuznogo soveshchaniya po spektroskopii. t. 1: Molekulyarnaya spektroskopiya (Papers of the 10th All-Union Conference on Spectroscopy. Vol. 1: Molecular Spectroscopy) [L'vov] Izd-vo L'vovskogo univ-ta, 1957. 499 p. 4,000 copies printed. (Series: Its: Fizichnyy zbirnyk, vyp. 3/8/)

Additional Sponsoring Agency: Akademiya nauk SSSR. Komissiya po spektroskopii. Ed.: Jaser, S.L.; Tech. Ed.: Saranyuk, T.V.; Editorial Board: Landberg, G.S., Academician (Resp. Ed., Deceased), Neporent, B.S., Doctor of Physical and Mathematical Sciences, Pabelinskiy, I.L., Doctor of Physical and Mathematical Sciences, Fabrikant, V.A., Doctor of Physical and Mathematical Sciences, Kornitskiy, V.G., Candidate of Technical Sciences, Rayskiy, S.M., Candidate of Physical and Mathematical Sciences, Klimovskiy, L.K., Candidate of Physical and Mathematical Sciences, Miliyanchuk, V.S., Candidate of Physical and Mathematical Sciences, and Glauberman, A. Ye., Candidate of Physical and Mathematical Sciences.

Card 1/30

| | |
|--|-----|
| Kolesova, V.A. Vibrational Spectra of Double-component Phosphate Glasses and Some Crystalline Phosphates | 461 |
| Mal'tsev, A.A., Ye. N. Moskritina, and V.M. Tatevskiy. Study of the Isotopic Effect and Verification of the Infrared Spectrum of Boron Trifluoride | 465 |
| Mal'tsev, A.A., Ye. N. Moskritina, and V.M. Tatevskiy. Quantitative Analysis of Boron Isotopes by Means of Infrared Spectra of Boron Trifluorides | 472 |
| Mal'tsev, A.A., Yu. Ya. Kuzakov, and V.M. Tatevskiy. Study of Electron Spectra and Isotopic Effect in Boron Oxygen Compounds | 475 |
| Mal'tsev, A.M., V.G. Vinokurov, and V.M. Tatevskiy. Study of Electron Spectra and Isotopic Effect in Boron Oxygen Compounds | 480 |

Card 29/30

TATEVSKIY, V.VI.

USSR / Chemistry of High Molecular Compounds.

L.

'Abs Jour : Ref. Zhur. - Khimiya, No.2, 1958, 6761.

Author : Salimov, H.A., Postovskaya, A.F., Kuz'minskiy, A.S.,
Tatevskiy, V.M.

Inst : Moscow University

Title : Investigation of Structural Changes of Sodium Butadiene
Caoutchouc in the Thermal Oxidation Process using Infra-
Red Spectroscopy Method.

Orig Pub : Vesti. Mosk. un-ta. Ser. matem., mekhan., astron., fiz.,
khimiya, 1957, No.1, 164-169.

Abstract : By studying the IK (infra-red) spectra of thermally ox-
idized sodium butadiene caoutchouc at 140^o (60 min. oxidation),
it has been established that during the oxidation process
the following oxygen-containing groups were formed:

Card : 1/8

USSR / Chemistry of High Molecular Compounds.

L.

Abs Jour : Ref. Zhur. - Khimiya, No.2, 1958, 6761.

Abstract : hydroperoxides, alcohols, esters, ethers, aldehydes, ketones and acids. After 20 minutes of oxidation there appeared an absorption band of -OOH and -OH groups at 3380 cm^{-1} the intensity of which increased with time of oxidation, while the maximum of absorption shifted to 3450 cm^{-1} (after 60 min.); the intensity of the absorption band of the carbonyl groups $\text{C}=\text{O}$ at 1700 cm^{-1} increased, while in the range of 1000-1400 cm^{-1} there appeared absorption bands of the groups $\text{R}_1-\text{C}(=\text{O})-\text{OR}$ at 1240 cm^{-1} and of the $\text{R}_1-\text{O}-\text{R}$ (ethers) at 1165 cm^{-1} . It was established that the newly appearing absorption bands at 810 and 880 cm^{-1} belong not to the peroxide, as it has been previously supposed, but to olefins having a structure 1,1,2-trialkylethylene $\text{R}_1\text{R}_2\text{C}=\text{CHR}_3$ and 1,1 dialkylethylene $\text{R}_1\text{R}_2\text{C}=\text{CH}_2$. In the course of oxidation, the intensity of the absorption bands of double bonds $\text{C}=\text{C}$ 1,2 (910 and 992 cm^{-1}) 1,4 - trans (965 cm^{-1}) and 1,4 - cis (680 cm^{-1}) was significantly decreased; an

Card : 2/3

USSR / Chemistry of High Molecular Compounds.

L.

Abs Jour : Ref. Zhur. - Khimiya, No.2, 1958, 6761.

Abstract : analogous decrease of intensities was also observed for absorption bands of double bonds C = C at 1640 cm^{-1} and for valence oscillation C-H of double bonds at 2979, 3019 and 3080 cm^{-1} ; a decrease of absorption bands of methylene groups -CH at 720, 2856 and 2929 cm^{-1} showed that O atoms in addition to double bonding attach themselves also to C atoms located in α position to the double bonds. An increase in intensity of the absorption band at 3400 cm^{-1} is linked to the formation of hydroxyl rather than hydroperoxide groups as it was confirmed by the results of the iodometric method.

Card : 3/3

The Apparatus for Production of Raman Scattering Spectra of
Gaseous Substances.

51-6-22/26

(Fig.1). The apparatus was used to study CO₂ gas. Spectrum of CO₂ at 3-6 atm was photographed using a spectrograph ИСН -51 with a short-focus camera. At 20 A in the light source the CO₂ doublet 1286-1388 cm⁻¹ was obtained after an exposure of several minutes. Fig. 2 shows this doublet obtained at 3 atm and 20 minute exposure. Two students of the Moscow State University, A.S. Bayshev and Yu.I. Kotov took part in this work. There are 2 figures and 5 references, 2 of which are Slavic.

SUBMITTED: January 12, 1957.

AVAILABLE: Library of Congress.

Card 2/2

TATEVSKIY, V. M.

51-3-3/14

AUTHORS: Pozdyshev, V. A., Pentin, Yu. A. and Tatevskiy, V. M.

TITLE: Infrared Spectra of Absorption by Certain Halogenated Alkanes in Liquid and Solid States and the Problem of Number and Configuration of Rotational Isomers.
(Infrakrasnyye spektry pogloshcheniya nekotorykh galoidalkanov v zhidkom i tverdom sostoyaniyakh i vopros o chisle i konfiguratsiyakh povorotnykh izomerov.)

PERIODICAL: Optika i Spektroskopiya, 1957, Vol.III, Nr.3, pp.211-220. (USSR)

ABSTRACT: At low temperatures concentrations of certain isomers are so small that their spectral lines or bands disappear. A "freezing" method based on this effect (Refs.1, 2, 3) was used by the present authors for comparison of infrared spectra of absorption by liquids and frozen substances. The results obtained were used to find the number, configuration and relative stability of rotational isomers in certain fairly complex halogenated alkanes.
Card 1/4 The following substances were studied: (1) 1,2-dichloroethane,

51-8-3/14

Infrared Spectra of Absorption by Certain Halogenated Alkanes in Liquid and Solid States and the Problem of Number and Configuration of Rotational Isomers.

(2) propyl chloride, (3) 1,2-dibromopropane, (4) 1,2-dibromobutane, (5) isobutyl bromide, (6) isoamyl bromide, (7) 1,4-dibromobutane. Infrared absorption spectra of these substances were obtained for the region of 450 to 1500 cm^{-1} . In freezing of the liquids the lowest temperature used was about -175°C . Figs. 2 & 3 show spectra of the substances studied in liquid and solid state respectively. For 1,2-dichloroethane three C_{2h} rotational forms were found and two C_2 forms. In the solid state only one C_{2h} form remains. Rotational isomeric configurations of propylchloride are two in number (C_3 and C_1 forms). In the solid state only one trans-form remains in propylchloride. The 618 cm^{-1} frequency was difficult to interpret. Infrared spectra of the liquid and crystalline states of isobutyl bromide were identical. The authors conclude that either C_3 and C_1 configurations exist both in the

Card 2/4

51-3-3/14

Infrared Spectra of Absorption by Certain Halogenated Alkanes in Liquid and Solid States and the Problem of Number and Configuration of Rotational Isomers.

liquid and solid states, or only one configuration exists in both states. The authors find themselves unable to decide this point. For 1,2-dibromopropane three rotational isomers were suggested. Again no difference between the liquid and solid-state spectra was observed. The 1138 cm^{-1} frequency, which is more intense in the solid state, is assigned to the trans-distribution of bromine atoms, the 1228 and 1203 cm^{-1} frequencies are assigned to the two other isomers. Molecules of isocamyl bromide and 1,2-dibromobutane have each two C-C bonds which permit rotational isomerism. In both these substances two rotational isomeric configurations are observed, both of which are stable in liquid and solid states. For 1,4-dibromobutane six theoretical configurations of the liquid state are shown in Fig.5. In the crystalline state of this substance only one isomer remains which has a mirror-rotational axis of the

Card 3/4

51-3-3/14

Infrared Spectra of Absorption by Certain Halogenated Alkanes in Liquid and Solid States and the Problem of Number and Configuration of Rotational Isomers.

second order (Fig. 5, Formula 6). There are 5 figures and 20 references, 4 of which are Slavic.

SUBMITTED: January 8, 1957.

AVAILABLE: Library of Congress

Card 4/4

POSTOVSKAYA, A.F.; SALIMOV, M.A.; KUZ'MINSKIY, A.S.; TATEVSKIY, V.M.

Structural changes of sodium-butadiene rubber in the process of
light oxidation. Fiz. sbor. no.3:423-427 '57. (MIRA 11:8)

1. Nauchno-issledovatel'skiy institut rezinovoy promyshlennosti i
Khimicheskii fakul'tet Moskovskogo ordena Lenina i ordena Trudovogo
Krasnogo Znameni gosudarstvennogo universiteta im. M.V. Lomonosova.
(Rubber---Spectra) (Oxidation)

TATEVSKIY, V.M.

Statistical calculations of the isotope exchange equilibria in the
gaseous state. Probl. kin. i kat. 9:339-344 '57. (MIRA 11:3)
(Gases) (Isotopes)

83635

S/081/60/000/015/001/014
A006/A001

5.2400A

Translation from: Referativnyy zhurnal, Khimiya, 1960, No. 15, p. 15, # 60232

AUTHORS: Mal'tsev, A.A., Kuzyakov, Yu.Ya., Tatevskiy, V.M. (I)
Mal'tsev, A.A., Vinokurov, V.G., Tatevskiy, V.M. (II)

TITLE: Study of Electron Spectra and of the Isotopic Effect in Oxygen
Boron Compounds. I. β -Bands of BO Molecules. II. "Boric Acid"
Bands

PERIODICAL: Fiz. sb. L'vovsk. un-t, 1957, No. 3 (8), pp. 475-480; 480-485

TEXT: I. A ДФC-3 (DFS-3) spectrograph (2A/mm dispersion) was used to investigate the emission spectrum of BO β -bands ($B^2\Sigma - X^2\Sigma$ transition) in the arc and a discharge tube with a hot hollow cathode containing B_2O_3 . Rotation analysis of 0 - 0, 0 - 1, 0 - 2, 0 - 3, 1 - 4, 1 - 5, 2 - 5, 2 - 6, and 3 - 4 bands was made, and by the method of least squares the following rotational constants (in cm^{-1}) of the $B^2\Sigma$ state were obtained: $B_e = 1.5192$, $\alpha_e = 0.0210$, $D_e = 7.4 \cdot 10^{-6}$ and $\beta_e = 2.0 \cdot 10^{-6}$. It is shown that divergence of Sheibe's rotational constant values (Sheibe, Z. Phys., 1930, Vol. 60, p. 74) with those of Djenkins and McKellar (Djenkins, McKellar, Phys. Rev. 1932, Vol. 42, p. 464)

Card 1/3

83635

S/081/60/000/015/001/014
A006/A001

Study of Electron Spectra and of the Isotopic Effect in Oxygen Boron Compounds.
I. β -Bands of BO Molecules. II. "Boric Acid" Bands

can be explained by the inaccurate treating of experimental data by Sheibe. The method of least squares was used to recalculate Sheibe's data for the $X^2\Sigma$ state. In all bands spin doubling was observed.

II. Spectrographs with diffraction gratings were used to investigate so-called fluctuation bands of boric acid, located in the 3700 - 6800 A range. The following spectrum sources were used: a discharge tube with a hot hollow cathode containing boron or boron-anhydride in an atmosphere of He and O₂ mixture, and an oxygen-hydrogen flame into which boric acid solution was introduced. At a high resolution the complicated rotational structure with several edges was observed for the majority of bands. The use of boron concentrated to 85% with a B¹⁰ isotope, allowed the determination of isotope band edges, shifted towards the short-wave side by about 6,5 and 5 A respectively for bands in the 5450 and 5750 A range. This result rejects Singh's theory (Singh, N.L., Proc. Indian Acad. Sci., 1949, Vol. A 29, p. 424) who relates the fluctuation bands of boric acid to the BO molecule. According to Singh the isotopic bands must be shifted to the long-wave side by 22 and 44 A respectively. When introducing to the spectrum source heavy water vapors, no isotopic effect is revealed in the

Card 2/3

83635

S/081/60/000/015/001/014
A006/A001

Study of Electron Spectra and of the Isotopic Effect in Oxygen Boron Compounds,
I. β -Bands of BO Molecules, II. "Boric Acid" Bands

fluctuation bands of the boric acid. This indicates the absence of hydrogen
in the molecule composition giving rise to these bands. It is assumed that the
fluctuation bands of the boric acid belong to the multi-atomic oxygen compound
of boron, B_xO_y .

A. Mal'tsev

Translator's note: This is the full translation of the original Russian
abstract.

X

Card 3/3

Chem. 115-1114
CH₂Cl, CH₂Br, CH₂I, CH₂F, CH₂Cl, CH₂Br, CH₂I, CH₂F, X
Cl, Br, I) were detd. The C-Br distance in CBr₄
is 1.93 Å; in the Br deriv. of ethylene it is 1.88 Å. The
distance C-I in the I₂ is 2.34 Å instead of 2.32 Å. cf.

TATAYEVSKY, Y. M.

7
1950